



Chylolymphatic Cyst with Midgut Volvulus and Malrotation in Pediatric Population: A Rare Triad and Comprehensive Scoping Review

Thangavel Prabhakar^{1*}, Himanshu Acharya², Vikesh Agrawal², Abhishek Tiwari², Rajpal Singh Sisodiya² and Dhananjaya Sharma¹

¹Department of General Surgery, NSCB Medical College and Hospital Jabalpur, Madhya Pradesh, India

²Department of Pediatric Surgery, NSCB Medical College and Hospital Jabalpur, Madhya Pradesh, India

Abstract

Introduction: Occurrence of chylolymphatic cyst with malrotation and midgut volvulus is a rare clinical entity and occurs as a result of lymphatic proliferation due to lymphatic obstruction caused by volvulus.

Methods: Since its first inception, very few studies have been published on this topic. We did a scoping review to identify the existence of the said condition in literature. We conducted this analysis using Mesh search terms "Midgut Volvulus", "Mesenteric Cyst", "Chylolymphatic Cyst" and "Malrotation" in PubMed, from the inception till current date.

Results: Scoping review revealed that only 13 cases are reported in literature so far. The cyst predominantly occurred in the jejunum (61%) and, to a lesser extent, in the ileum (30%) among the cases investigated. Most postoperative recoveries proceeded without complications, barring two cases identified during postmortem examinations. Intraoperative findings included the presence of a Ladd's band in 5 out of the 13 cases; however, the referenced studies did not note significant additional anomalies, except for congenital atresia of the transverse colon.

Conclusion: As this is a rare combination of two congenital anomalies, preoperative clinical diagnosis is difficult and requires careful evaluation and surgical management.

Keywords: Chylolymphatic Cyst; Midgut Volvulus; Malrotation.

INTRODUCTION

Malrotation of the midgut is a congenital anomaly originating during embryonic development, characterized by improper rotational alignment around the axis of the superior mesenteric artery. This condition leads to bilious emesis and acute intestinal obstruction [1]. Midgut volvulus represents a medical emergency with the potential to progress to bowel ischemia, necrosis, and, in severe cases, bowel perforation. It necessitates prompt surgical intervention, as no effective nonsurgical treatments currently exist [2]. Mesenteric chylolymphatic cysts, rare congenital lesions occurring in approximately 1 in 20,000 cases in the pediatric population, are thought to arise due to the benign proliferation of ectopic lymphatic tissue within the mesentery due to lymphatic obstruction resulting from the twist of the volvulus [3]. Mesenteric cysts are rare intra-abdominal masses in the pediatric population, with the chylolymphatic variant comprising only 7.3% of all abdominal cysts [3]. The co-occurrence of these anomalies is even rarer and is sparsely documented in existing literature. We conducted a rapid scoping review of previous studies to confirm its rarity.

Submitted: 01 April, 2024 | **Accepted:** 14 May, 2024 | **Published:** 16 May, 2024

***Corresponding author:** Thangavel Prabhakar, NSCB Medical College and Hospital, Jabalpur, Madhya Pradesh, India

Copyright: © 2024 Prabhakar T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Prabhakar T, Acharya H, Agrawal V, Tiwari A, Sisodiya RS, et al. (2024) Chylolymphatic Cyst with Midgut Volvulus and Malrotation in Pediatric Population: A Rare Triad and Comprehensive Scoping Review. SM J Pediatr Surg 7: 4.

MATERIALS AND METHODS

As this anomaly is sparsely reported, we conducted a scoping review and identified 8 case reports and 1 case series available as of October 2023, all dating back to the inception of the literature search [3, 8-14] (Figure 1, Table 1), with Inclusion criteria as pediatric patients less than 18 years having mesenteric cyst, mid gut malrotation, without any other congenital anomalies or iatrogenic/ traumatic mesenteric cysts. Overall, 13 cases fulfilling the inclusion criteria were taken for review, for which the PRISMA diagram is shown below. We conducted this analysis using Mesh search terms "Midgut Volvulus", "Mesenteric Cyst", "Chylolymphatic Cyst" and "Malrotation". In PubMed, from the inception until current date.

RESULTS

Notably, an almost equal incidence of chylolymphatic cysts was observed in both males and females. Commonly reported symptoms included abdominal pain, distension, and previous episodes. However, abdominal examination revealed mesenteric cysts in only 3 out of the 13 cases (23%). Imaging techniques, such as ultrasound and computed tomography, identified mid-gut malrotation with volvulus in 5 cases (38% sensitivity), with 3 cases showing cysts alone. Further literature review indicated that the cyst was predominantly located in the jejunum (61%) and the ileum (30%). Postoperative recoveries were generally uneventful, except for 2 cases diagnosed postmortem. Intraoperatively, a Ladd's band was observed in 5 out of the 13 cases, and the mentioned studies did not report significant additional anomalies, except for congenital atresia of the transverse colon found in a study by Fukuta A, Inoue T, Kawakubo N, Kondo T, Matsuura T [15].

Discussion and Scoping Review

Midgut volvulus, a congenital rotational anomaly, arises due to non-rotation, reverse rotation, partial rotation, or adherence to the mesentery following the physiological herniation of bowel loops around the superior mesenteric artery (SMA) axis during the embryological period [1]. The Ladd's procedure is performed by open or laparoscopic techniques to




Figure 1

Table 1: Table showing observations of rapid scoping review of literature on mesenteric cyst associated with midgutmalrotation.

S. No	Author	Year	Age (yrs)	Gender	Clinical identification	Identification on Imaging	Location	Surgery Done
1	KronSD [8]	1954	6	Male	No	No	Jejunum	Excision
2	Bentley JFR [9]	1959	6	Female	Yes	No	Jejunum	Intestinal Resection
3	Bentley JFR [9]	1959	4	Female	No	No	Ileum	Intestinal Resection
4	Bentley JFR [9]	1959	6	Female	No	No	Jejunum	IS(death)
5	Bentley JFR [9]	1959	2	Male	No	0	Ileum	IS(death)
6	Namasivayam J[10]	1992	10	Female	Yes	Yes	Jejunum	Excision
7	Weeda B [11]	2008	6	Male	No	Yes	Jejunum	Intestinal Resection
8	Weeda B [11]	2008	0	Male	No	Yes	Jejunum	Intestinal Resection
9	Botchway E [12]	2012	2	Male	No	Yes	Present	Excision
10	Botchway E [12]	2012	9	Male	No	Yes	Ileum	Intestinal Resection
11	Singh J [13]	2013	22	Male	No	Yes	Ileum	Excision
12	Alfadhel SF [14]	2019	2	Male	No	Yes	Ileum	Resection
13	Pai PK [3]	2021	16	Female	Yes	Yes	Jejunum	Excision

IS- in-situ, couldn't be operated.



reduce volvulus and remove Ladd bands if present [4]. The derotation of volvulus is relatively easy by open technique and involves anticlockwise derotation which can be performed using “steering wheel technique” in laparoscopy [5]. Post-surgical recurrence risk is reduced by widening the mesentery, with recurrence rates ranging from 2% to 8% [6]. Bowel loops are examined for necrosis, a potential cause of short bowel syndrome in cases of acute midgut volvulus [7]. In the present case, it was complicated by a chylolymphatic cyst, which needed resection.

Notably, we observed an almost equal incidence between males and females in these cases. Commonly reported symptoms across most presentations included abdominal pain, abdominal distension, and a previous episode. However, abdominal examination findings indicated mesenteric cysts in only 3 out of the 13 cases (23%). Imaging techniques, including ultrasound and computed tomography, identified mid-gut malrotation with volvulus in only 5 out of the 13 cases (38% sensitivity), with 3 cases showing cysts alone. Further review of literature reveals, that the cyst was most frequently located in the jejunum (61%), followed by the ileum (30%) in the cases examined. Postoperative recoveries were generally uneventful, except for 2 cases that were diagnosed after conducting postmortem. A Ladd’s band was observed intraoperatively in 5 out of the 13 cases, however, the aforementioned studies did not report significant additional anomalies, with the exception of congenital atresia of the transverse colon found in a study by Fukuta A, Inoue T, Kawakubo N, Kondo T, Matsuura T [15].

Mesenteric cysts are fluid-filled structures located within the mesentery, and they may or may not extend into the retroperitoneum, often featuring a mesothelial cell lining [16]. These cysts vary in size, ranging from small sub-centimetric cysts to larger ones that resemble complex ascites [6]. In the present case, intraoperative findings revealed a dumbbell-shaped fluctuant cystic lesion partially enveloping the jejunal loop, lacking major feeding vessels. Small mesenteric cysts are challenging to diagnose clinically, and provisional diagnosis is typically reliant on imaging, which can miss such cyst. Recurrence and malignancy are rare in this condition, although secondary complications may include fluid leakage leading to infection, bowel loop herniations, and obstructions [17]. A confirmed diagnosis is typically established through laparotomy and histopathological assessment. However, it could not be confirmed if the mesenteric cyst was causing the volvulus or if both entities are a mere coexistence.

CONCLUSION

In conclusion, midgut volvulus, a congenital rotational anomaly, presents a complex scenario in pediatric surgery. The presence of chylolymphatic cyst associated in these cases, while uncommon, adds an intriguing dimension to the understanding of midgut volvulus. Their diagnostic challenges and rarity emphasize the clinical importance of identification and management of said condition. A scoping review in the manuscript adds to the information on this particular entity which confirms the rarity of the condition.

ETHICAL APPROVAL

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

AUTHOR CONTRIBUTIONS

This is to declare that all authors have contributed to the study. No part of the manuscript has been sent for consideration elsewhere or published in any International or National journal. The authors clearly certify that there is no aspect of plagiarism. All the conflicts of interest have been clearly defined and the source of grant disclosed. Due ethical permission/consent has been obtained for carrying out the study. In case

of any dispute, the authors will be held fully responsible for the statement disclosed in the cover letter. The authors are also aware of the copyright rules and also declare that they will not reproduce any published text without due permission from the journal.

REFERENCES

1. Alani M, Rentea RM. Midgut Malrotation. [Updated 2023 Jul 31]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024.
2. Pandey A, Singh SP, Gupta V, Pandey J, Sachan P. Malrotation with midgut volvulus associated with perforated ileal duplication. *J Indian Assoc Pediatr Surg.* 2013; 18(4): 155-157. doi: 10.4103/0971-9261.121123. PMID: 24347871; PMCID: PMC3853859.
3. Pai PK, Gudmundsdottir H, Hull NC, Thacker PG, Klinkner DB. Chylolymphatic mesenteric cyst with midgut volvulus in an adolescent: a peculiar presentation. *Radiol Case Rep.* 2021; 17(2): 420-422. doi: 10.1016/j.radcr.2021.10.066. Erratum in: *Radiol Case Rep.* 2023; 18(4): 1641-1642. PMID: 34934468; PMCID: PMC8654696.
4. Xie W, Li Z, Wang Q, Wang L, Pan Y, Lu C. Laparoscopic vs open Ladd’s procedure for malrotation in neonates and infants: a propensity score matching analysis. *BMC Surg [Internet].* 2022; 22.
5. Agrawal V, Tiwari A, Acharya H, Mishra R, Sharma D. Laparoscopic ‘steering wheel’ derotation technique for midgut volvulus in children with intestinal malrotation. *J Minim Access Surg.* 2019; 15(3): 219-223. doi: 10.4103/jmas.JMAS_24_18. PMID: 29737312; PMCID: PMC6561056.
6. Zamir D, Yuchtman M, Amar M, Shoemo U, Weiner P. [Giant mesenteric cyst mimicking ascites]. *Harefuah.* 1996; 130(10): 683-684, 727. Hebrew. PMID: 8794659.
7. Svetanoff WJ, Sobrino JA, Sujka JA, St Peter SD, Fraser JD. Laparoscopic Ladd Procedure for the Management of Malrotation and Volvulus. *J Laparoendosc Adv Surg Tech A.* 2020; 30(2): 210-215. doi: 10.1089/lap.2019.0602. Epub 2019 Dec 31. PMID: 31895626.
8. KRON SD, SATINSKY VP. Chylous mesenteric cyst causing volvulus of malrotated intestine. *AMA Am J Dis Child.* 1954; 88(2): 216-279. doi: 10.1001/archpedi.1954.02050100218009. PMID: 13180016.
9. BENTLEY JF, O’DONNELL MB. Mesenteric cysts with malrotated intestine. *Br Med J.* 1959; 2(5146): 223-225. doi: 10.1136/bmj.2.5146.223. PMID: 13671235; PMCID: PMC1990025.
10. Namasivayam J, Ziervogel MA, Hollman AS. Case report: volvulus of a mesenteric cyst—an unusual complication diagnosed by CT. *Clin Radiol.* 1992; 46(3): 211-212. doi: 10.1016/s0009-9260(05)80450-7. PMID: 1395431.
11. Weeda VB, Booij KA, Aronson DC. Mesenteric cystic lymphangioma: a congenital and an acquired anomaly? Two cases and a review of the literature. *J Pediatr Surg.* 2008; 43(6): 1206-1208. doi: 10.1016/j.jpedsurg.2008.01.075. PMID: 18558209.
12. Botchway E, Tshifularo N, Human MJ. Mesenteric cyst as a presentation of chronic midgut volvulus in a 12-year-old boy. *J Pediatr Surg Case Rep.* 2020; 55(101351): 101351.
13. Singh N, Singh R, Maheswari U, Aga P. Primary mesenteric lymphangioma in a young adult with intestinal malrotation and ‘counter-clockwise barber pole sign’. *BMJ Case Rep.* 2013; 2013:bcr2013008994. doi: 10.1136/bcr-2013-008994. PMID: 23761495; PMCID: PMC3702800.



14. Alfadhel SF, Alghamdi AA, Alzahrani SA. Ileal volvulus secondary to cystic lymphangioma: A rare case report with a literature review. *Avicenna J Med.* 2019; 9(2): 82-85. doi: 10.4103/ajm.AJM_203_18. PMID: 31143702; PMCID: PMC6530267.
15. Fukuta A, Inoue T, Kawakubo N, Kondo T, Matsuura T. Congenital intestinal atresia associated with a mesenteric cystic lymphangioma in a low birth weight neonate: A case report. *Int J Surg Case Rep.* 2020; 75: 136-139. doi: 10.1016/j.ijscr.2020.09.021. Epub 2020 Sep 10. PMID: 32949911; PMCID: PMC7502782.
16. Kurtz RJ, Heimann TM, Holt J, Beck AR. Mesenteric and retroperitoneal cysts. *Ann Surg.* 1986; 203(1): 109-112. doi: 10.1097/0000658-198601000-00017. PMID: 3942415; PMCID: PMC1251046.
17. Hassan M, Dobrilovic N, Korelitz J. Large gastric mesenteric cyst: case report and literature review. *Am Surg.* 2005; 71(7): 571-573. PMID: 16089120.